#76

NOAA Technical Report NMFS 76

April 1989

Analysis of Fish Diversion Efficiency and Survivorship in the Fish Return System at San Onofre Nuclear Generating Station

Milton S. Love
Meenu Sandhu
Jeffrey Stein
Kevin T. Herbinson
Robert H. Moore
Michael Mullin
John S. Stephens, Jr.

LIBRARY

MAY 2 1989

National Marine Fisheries Service
2570 Dole Street
Honolulu, Hawaii

NOAA
Tech.
Rept.
NMFS
76

U.S. Department of Commerce

NOAA TECHNICAL REPORT NMFS

The major responsibilities of the National Marine Fisheries Service (NMFS) are to monitor and assess the abundance and geographic distribution of fishery resources, to understand and predict fluctuations in the quantity and distribution of these resources, and to establish levels for their optimum use. NMFS is also charged with the development and implementation of policies for managing national fishing grounds, development and enforcement of domestic fisheries regulations, surveillance of foreign fishing off United States coastal waters, and the development and enforcement of international fishery agreements and policies. NMFS also assists the fishing industry through marketing service and economic analysis programs, and mortgage insurance and vessel construction subsidies. It collects, analyzes, and publishes statistics on various phases of the industry.

The NOAA Technical Report NMFS series was established in 1983 to replace two subcategories of the Technical Reports series: "Special Scientific Report—Fisheries" and "Circular." The series contains the following types of reports: Scientific investigations that document long-term continuing programs of NMFS; intensive scientific reports on studies of restricted scope; papers on applied fishery problems; technical reports of general interest intended to aid conservation and management; reports that review in considerable detail and at a high technical level certain broad areas of research; and technical papers originating in economics studies and from management investigations. Since this is a formal series, all submitted papers receive peer review and those accepted receive professional editing before publication.

Copies of NOAA Technical Reports NMFS are available free in limited numbers to governmental agencies, both Federal and State. They are also available in exchange for other scientific and technical publications in the marine sciences. Individual copies may be obtained from: U.S. Department of Commerce, National Technical Information Service, 5285 Port Royal Road, Springfield, VA 22161. Although the contents have not been copyrighted and may be reprinted entirely, reference to source is appreciated.

34. Additions to a revision of the shark genus *Carcharhinus*: Synonymy of *Aprionodon* and *Hypoprion*, and description of a new species of *Carcharhinus* (Carcharhinidae), by J. A. F. Garrick. November 1985, 26 p.

35. Synoptic review of the literature on the Southern oyster drill *Thais haemastoma floridana*, by Philip A. Butler. November 1985, 9 p.

36. An egg production method for estimating spawning biomass of pelagic fish: Application to the northern anchovy, *Engraulis mordax*, by Reuben Lasker (editor). December 1985, 99 p.

37. A histopathologic evaluation of gross lesions excised from commercially important North Atlantic marine fishes, by Robert A. Murchelano, Linda Despres-Patanjo, and John Ziskowski. March 1986, 14 p.

38. Fishery atlas of the northwestern Hawaiian Islands, by Richard N. Uchida and James H. Uchiyama (editors). September 1986, 142 p.

39. Survey of fish protective facilities at water withdrawal sites on the Snake and Columbia Rivers, by George A. Swan, Tommy G. Withrow, and Donn L. Park. April 1986, 34 p.

40. Potential impact of ocean thermal energy conversion (OTEC) on fisheries, by Edward P. Myers, Donald E. Hoss, Walter M. Matsumoto, David S. Peters, Michael P. Seki, Richard N. Uchida, John D. Ditmars, and Robert A. Paddock. June 1986, 33 p.

41. A stationary visual census technique for quantitatively assessing community structure of coral reef fishes, by James A. Bohnsack and Scott P. Bannerot. July 1986, 15 p.

42. Effects of temperature on the biology of the northern shrimp, *Pandalus borealis*, in the Gulf of Maine, by Spencer Apollonio, David K. Stevenson, and Earl E. Dunton, Jr. September 1986, 22 p.

43. Environment and resources of seamounts in the North Pacific, by Richard N. Uchida, Sigeiti Hayasi, and George W. Boehlert (editors). September 1986, 105 p.

44. Synopsis of biological data on the porgies, *Calamus arctifrons* and *C. proridens* (Pisces: Sparidae), by George H. Darcy. September 1986, 19 p.

45. Meristic variation in *Sebastes* (Scorpaenidae), with an analysis of character association and bilateral pattern and their significance in species separation, by Lochai Chen. September 1986, 17 p.

46. Distribution and relative abundance of pelagic nonsalmonid nekton off Oregon and Washington 1979-84, by Richard D. Brodeur and William G. Pearcy. December 1986, 85 p.

47. Reproduction, maturation, and seed production of cultured species: Proceedings of the twelfth U.S.-Japan meeting on aquaculture, Baton Rouge, Louisiana, October 25-29, 1983, by Carl J. Sindermann (editor). February 1987, 73 p.

48. Widow rockfish: Proceedings of a workshop, Tiburon, California, December 11-12, 1980, by William H. Lenarz and Donald R. Gunderson (editors). January 1987, 57 p.

49. Reproduction, movements, and population dynamics of the southern kingfish, *Menticirrhus americanus*, in the northwestern Gulf of Mexico, by Stephen M. Harding and Mark E. Chittenden, Jr. March 1987, 21 p.

50. Preparation of acetate peels of valves from the ocean quahog, *Arctica islandica*, for age determinations, by John W. Ropes. March 1987, 5 p.

51. Status, biology, and ecology of fur seals: Proceedings of an international workshop, Cambridge, England, 23-27 April 1984, by John P. Croxall and Roger L. Gentry (editors). June 1987, 212 p.

52. Limited access alternatives for the Pacific groundfish fishery, by Daniel D. Huppert (editor). May 1987, 45 p.

53. Ecology of east Florida sea turtles: Proceedings of the Cape Canaveral, Florida, sea turtle workshop, Miami, Florida, February 26-27, 1985, by Wayne N. Witzell (convener and editor). May 1987, 80 p.

54. Proximate and fatty acid composition of 40 southeastern U.S. finfish species, by Janet A. Gooch, Malcolm B. Hale, Thomas Brown, Jr., James C. Bonnet, Cheryl G. Brand, and Lloyd W. Reiger. June 1987, 23 p.

55. Proximate composition, energy, fatty acid, sodium, and cholesterol content of finfish, shellfish, and their products, by Judith Krzynowek and Jenny Murphy. July 1987, 53 p.

56. Some aspects of the ecology of the leatherback turtle *Dermochelys coriacea* at Laguna Jolova, Costa Rica, by Harold F. Hirth and Larry H. Ogren. July 1987, 14 p.

57. Food habits and dietary variability of pelagic nekton off Oregon and Washington, 1979-1984, by Richard D. Brodeur, Harriet V. Lorz, and William G. Pearcy. July 1987, 32 p.

58. Stock assessment of the Gulf menhaden, *Brevoortia patronus*, fishery, by Douglas S. Vaughan. September 1987, 18 p.

59. Atlantic menhaden, *Brevoortia tyrannus*, purse seine fishery, 1972-84, with a brief discussion of age and size composition of the landings, by Joseph W. Smith, William R. Nicholson, Douglas S. Vaughan, Donnie L. Dudley, and Ethel A. Hall. September 1987, 23 p.

60. Gulf menhaden, *Brevoortia patronus*, purse seine fishery, 1974-85, with a brief discussion of age and size composition of the landings, by Joseph W. Smith, Eldon J. Levi, Douglas S. Vaughan, and Ethen A. Hall. December 1987, 8 p.

61. Manual for starch gel electrophoresis: A method for the detection of genetic variation, by Paul B. Aebersold, Gary A. Winans, David J. Teel, George B. Milner, and Fred M. Utter. December 1987, 19 p.

62. Fishery publication index, 1980-85; Technical memoradum index, 1972-85, by Cynthia S. Martin, Shelley E. Arenas, Jacki A. Guffey, and Joni M. Packard. December 1987, 149 p.

63. Stock assessment of the Atlantic menhaden, *Brevoortia tyrannus*, fishery, by Douglas S. Vaughan and Joseph W. Smith. January 1988, 18 p.

64. Illustrated key to penaeoid shrimps of commerce in the Americas, by Isabel Pérez Farfante. April 1988, 32 p.

65. History of whaling in and near North Carolina, by Randall R. Reeves and Edward Mitchell. March 1988, 28 p.

66. Atlas and zoogeography of common fishes in the Bering Sea and northeastern Pacific, by M. James Allen and Gary B. Smith. April 1988, 151 p.

NMFS Honolulu Library



NOAA Technical Report NMFS 76

Analysis of Fish Diversion Efficiency and Survivorship in the Fish Return System at San Onofre Nuclear Generating Station

Milton S. Love
Meenu Sandhu
Jeffrey Stein
Kevin T. Herbinson
Robert H. Moore
Michael Mullin
John S. Stephens, Jr.

April 1989

U.S. DEPARTMENT OF COMMERCE
Robert Mosbacher, Secretary
National Oceanic and Atmospheric Administration
William E. Evans, Under Secretary for Oceans and Atmosphere
National Marine Fisheries Service
James Brennan, Assistant Administrator for Fisheries

Library
NOAA, National Marine Fisheries Service
SWFC Honolulu Laboratory F/SWC2
2570 Dole Street
Honolulu, HI 96822-2396

The National Marine Fisheries Service (NMFS) does not approve, recommend or endorse any proprietary product or proprietary material mentioned in this publication. No reference shall be made to NMFS, or to this publication furnished by NMFS, in any advertising or sales promotion which would indicate or imply that NMFS approves, recommends or endorses any proprietary product or proprietary material mentioned herein, or which has as its purpose an intent to cause directly or indirectly the advertised product to be used or purchased because of this NMFS publication.

Contents

Introduction 1
Description of SONGS Units 2 and 3 1
Description of the fish return system 2

Methods and materials 2
Diversion efficiency 2
Survivorship 4
Test experiments 4
Reference experiments 5

Results 6
Diversion efficiency 6
Survivorship 9

Discussion 10

Acknowledgments 12

Citations 13

Appendices 14



Digitized by the Internet Archive
in 2025

https://archive.org/details/noaa-technical-report-nmfs-ssrf_1989-04_76

An Analysis of Fish Diversion Efficiency and Survivorship in the Fish Return System at San Onofre Nuclear Generating Station

MILTON S. LOVE[1]
JEFFREY STEIN[2]
ROBERT H. MOORE
MICHAEL MULLIN
JOHN S. STEPHENS, JR.
VANTUNA Research Group
Moore Laboratory of Zoology
Occidental College
Los Angeles, California 90041

MEENU SANDHU
Rand Corporation
1700 Main Street
Santa Monica, California 90406

KEVIN T. HERBINSON
Environmental Affairs
Southern California Edison
P.O. Box 800
Rosemead, California 91770

ABSTRACT

This study examined the efficiency of fish diversion and survivorship of diverted fishes in the San Onofre Nuclear Generating Station Fish Return System in 1984 and 1985. Generally, fishes were diverted back to the ocean with high frequency, particularly in 1984. Most species were diverted at rates of 80% or more. Over 90% of the most abundant species, *Engraulis mordax*, were diverted. The system worked particularly well for strong-swimming forms such as *Paralabrax clathratus*, *Atherinopsis californiensis*, and *Xenistius californiensis*, and did not appreciably divert weaker-swimming species such as *Porichthys notatus*, *Heterostichus rostratus*, and *Syngnathus* sp. Return rates of some species were not as high in 1985 as in 1984. Individuals of most tested species survived both transit through the fish return system and 96 hours in a holding net. Some species, such as *E. mordax*, *X. californiensis*, and *Umbrina roncador*, experienced little or no mortality. Survivorship of *Seriphus politus* was highly variable and no *Anchoa delicatissima* survived.

Introduction

Coastal steam-electric plants in southern California obtain cooling water from 900-1000 m offshore via pipes 3-4 m in internal diameter. Water enters a bowl-like intake and flows shoreward to the plant, where traveling screens act as strainers. The filtered water flows through condenser cooling tubes and is discharged back into the ocean.

Fish may enter into these intakes (Helvey 1985, 1987) and are pulled shoreward swimming against the current (Downs and Meddock 1974). Small fishes or weaker-swimming species are impinged on the traveling screens and removed for disposal; stronger individuals may reside in the screenwells in front of the screens for some time. The number of fishes killed in this manner can be large at times (Schuler and Larson 1975). Various control technologies have been developed to reduce fish loss at intake structures (EPRI 1981).

To minimize fish losses at the San Onofre Nuclear Generating Station (SONGS), Southern California Edison (SCE) has incorporated a fish return system (FRS) into the Units 2 and 3 cooling water system (Downs and Meddock 1974). In this system, fish may either be impinged on travelling screens or diverted into the fish return system. The system was, and remains, unique (Lawler et al. 1982a, EPRI 1984), and SCE contracted with the VANTUNA Research Group of Occidental College to determine its efficiency during 1984 and 1985. This report describes the results of that study. The objectives of this study were 1) to examine the efficiency of fish diversion within the SONGS return system, and 2) to test the survivorship of the diverted fish.

Description of SONGS Units 2 and 3

SONGS is located on the California coast, at 33°22′N and 117°33.5′W, between San Clemente and Oceanside (Fig. 1). SONGS Units 2 and 3 are 1,100 MW facilities. Unit 2 became commercially operational in August 1983, Unit 3 in April 1984. Each unit has a once-through cooling system, with flow rates of 3,137 m^3/min (830,000 gpm). The intakes are located 970.2 m offshore, at a depth of 9.8 m (Fig. 1). Both units have diffuser-type discharges, consisting of 63 ports spread over a distance of 750 m. The Unit 2 diffuser begins 1,795 m offshore extending to 2,545 m, at depths of ll.9-14.9 m. The Unit 3 diffuser begins 1,084 m offshore and extends to 1,830 m, at depths from 9.8 to 11.6 m (Yuge and Herbinson 1985).

Present addresses:
[1] Marine Science Institute, University of California at Santa Barbara, Santa Barbara, CA 93106.
[2]Marine Biology Research Division, Scripps Institution of Oceanography, University of California at San Diego, La Jolla, CA 92037.



Figure 1
Location of the San Onofre Nuclear Generating Station (SONGS) and its fish return system.

Description of the fish return system (FRS)

The FRS (Figs. 2,3) relies on behavioral responses of fish to varying water velocities and pressures. Fish within the cooling water encounter concrete vanes and angled plastic louvers situated in front of the traveling screens. These are angled toward a bypass area and create a pressure differential detected by the fish, which swim along the louvers (Downs and Meddock 1974, Schuler and Larson 1975). The bypass area, a quiet-water concrete-lined basin, measures 4.9 × 4 m. A watertight elevator basket, open at the top, sits within this basin. When activated, the elevator ascends, collecting most of the fish in the basin. Upon reaching its maximum height, the elevator tips, spilling the fish into a sluice channel. This procedure is repeated several times until most fish are removed. Simultaneously, additional water flushes into the channel and the fishes are discharged into a pipe which empties in 6 m of water, about 400 m offshore.

Methods and materials

Diversion efficiency

Twice-weekly impingement and elevator samples were taken at Units 2 and 3 to establish the diversion efficiency of the FRS. We attempted two 24-hr samples on consecutive days during the week (though this was not always possible during periods of variable plant operation). Twenty-four hrs before sampling, the screens were cleaned of impinged fishes and the FRS was run to remove all entrapped fishes. Diversion efficiency was calculated for fish from samples that conformed to the following criteria: 1) The sample was drawn from a 22-26 hr impingement or elevator sample; 2) impingement and elevator samples were taken on the same day; and 3) all four circulating pumps were operating at the sampled unit. A total of 55 samples at Unit 2 and 65 samples at Unit 3 conformed to these criteria in 1984 and 1985. Of these samples, 35 impingement and elevator samples were taken on the same day. Fish from these samples were evaluated for diversion efficiency by species and size.

On each sample day, the traveling screens were run to wash any impinged fish into the impingement basket. At the same time, the contents of the elevator basket were dumped into the return sluiceway. Using two 15-inch diameter nets, two aliquot samples (sampling about one-tenth of all fish) were taken as fish were poured from the basket. This procedure was repeated until fish were no longer collected in the basket (three to four repetitions). Fish collected from the elevator samples and the impingement basket were identified, counted, measured, and weighed according to criteria previously described (DeMartini et al. 1984).

Return efficiency was computed for all fishes that went through the fish return system. Of the fish returned, two species, the northern anchovy (*Engraulis mordax*) and queenfish (*Seriphus politus*), accounted for a large percentage of all individuals. Detailed analyses were conducted for the most abundant fish—*E. mordax*, *S. politus*, white croaker (*Genyonemus lineatus*), walleye surfperch (*Hyperprosopon argenteum*, and white surfperch (*Phanerodon furcatus*). Other species were not analyzed due to the lack of sufficient measurement data. An attempt to group all other species into one category was abandoned because of the wide range of behavior of the remaining species.

During a sample period, thousands of fish go through the fish return system of which only a small proportion get impinged. Out of the large numbers of fish collected during each sample, approximately 200-400 fish were randomly sampled and measured for size from the fish return system. Similarly, 200-400 fish were measured for the impinged samples. The latter, however, represented almost all of the fish in the sample. In order to get a representation of the number of fish in each size-class that were diverted throughout the system or impinged on the screens, we computed



Figure 2
Fish return system of San Onofre Units 2 and 3, from intake through fish removal area.


TRAVELING WATER SCREEN
FISH ELEVATOR DRAW WORKS DRIVE MECHANISM SUPPORT FRAMES AND GUIDES
FISH FLUSHING MANIFOLD
ELEVATION 16'-0"
LOCAL CONTROL CONSOLE
FISH ELEVATOR BUCKET
SLUICING WATER INLET LINE
FISH SLUICING DISCHARGE SEA LINE
OPEN SLUICING CHANNEL
FISH HOLDING CHAMBER (DASHED LINE PROFILE)
FLOW DIVIDING BARRIER (PART OF INTAKE STRUCTURE)
FISH BUCKET AT LOWEST POSITION EL.(-)26'-0" APPROX
FISH INLET FLOW 25' DEEP

Figure 3
Removal elevator and sluicing channel of fish return system, San Onofre Units 2 and 3.



Figure 4
Holding net used in test phase of the 96-hour survivorship studies.

the proportion within each size-class and extrapolated the proportions to the total number that went through the fish return system. These adjusted data are what were used in the analyses. The actual number of fish measured in each 10-mm size-class by year is presented in Appendix A.

Our analyses tested the null hypothesis that the return efficiency did not vary under different conditions or by fish size. The conditions tested were year (1984, 1985) and units (Units 2,3). We tested the return efficiency between years, sites, and fish size-class (<100 mm, >100 mm standard length), using the three-way analysis of variance. (Note that return efficiency figures in Figures 6-10 are plotted in 10-mm size-class increments. We feel that this is more explicative than showing only two size-classes.) The ANOVA design was unbalanced since species may not necessarily be impinged and/or returned at each sampling time or in each size-class. Multiple comparisions were also performed using the Bonferroni test (Bailey 1977). Natural log (1% return) transformation (referred to as "transformed return efficiency" throughout the text) was utilized to remove the skewness of the data and heterogeneity of variance. The variable (1% return) was used to change the negatively skewed distribution to a positively skewed distribution which closely approximated the log normal distribution. Although other transformations were tried, it was found that the above transformation was the best in upholding the homogeneity of variance assumption necessary for the analysis of variance.

Due to the natural sequential order of time in the data, autocorrelation may be present which would violate the assumption of sample independence. The data were checked for autocorrelation by using sinusoidal curve fit where the transformed return efficiency was regressed against time for the five species by size-class. The sinusoidal curve fit was used, since the samples were irregularly spaced in time (Lorda and Saila 1986).

Plots of percent return efficiency were done by size-class, site, and year to examine the results from the analysis of variance.

Survivorship

Test experiments We evaluated the survival rates of fish transported through the FRS by conducting *in situ* 96-hr survivorship tests at the terminus of the fish return conduit. Ninety-six hours is a standard assay period and represented the longest experimental time-sequence in which we felt animals could be maintained in holding nets without experimental bias. The experiments were on 24-hr samples of entrapped fish discharged offshore from the return conduit into a 3.5 m^2 holding net (Fig. 4). All survivorship results were obtained with a biologist coordinating the operation of the FRS in-plant. All phases of the operational procedure—running the sluicing water, lifting the elevator basket, and flushing the conduit—were done to minimize the trauma incurred by the returned fish.

Before connecting the net, we flushed the return conduit of any resident fish by running the conduit flushing water for 15 minutes. At the start of an experiment, an in-plant biologist coordinated the dumping of the fish elevator to coincide with offshore biologists connecting the holding net to the return conduit. After dumping the elevator contents, we flushed the conduit for an additional 15 min to insure that all fish were discharged into the net. At this point, the net was disconnected from the conduit and moved approximately 10 m either up or downcoast and secured to the bottom. Divers then assessed the number, condition, and species-composition of fish within the net and removed any dead individuals. Fish condition and behavior were assessed once every 24 hrs until the end of the test period, at which time all surviving fish were identified, counted, measured, and



Figure 5
Holding net used in the 96-hour reference studies.

inspected for lesions and abrasions derived from the flushing process. Predation within the net was minimized by removing piscivorous species, such as the kelp bass (*Paralabrax clathratus*).

Reference experiments To evaluate stress caused by containment within the nets, free-swimming fish were captured and held within the holding nets for 96-hr reference experiments. The reference specimens were captured by fishing the holding net with fyke wings (Fig. 5), thereby allowing fish to enter the net without the stress associated with being discharged through the return conduit. The reference tests were designed to determine holding-net effects on survivorship.

The reference nets were set with fyke wings offshore of San Onofre in the vicinity of the fish return conduit. After allowing a net to fish for 24 hrs, we disconnected the wings and sealed the net if sufficient numbers of fish were captured. We set the net one day prior to the start of the test. This method allowed the start of both experiments to coincide. As with the test nets, the reference nets were inspected every 24 hrs, and at the end of the 96-hr holding period all fish were collected and the appropriate data recorded.

We tested the null hypothesis that there was no difference in fish survivorship between site, years, and fish length. Comparisons were made between reference and return survivorship. Due to limited numbers of fish sampled, *E. mordax*, *G. lineatus*, and *S. politus* could be used in the analyses. Plots presenting the conditional probability of survival by each day and size (over the 96 hr-period) suggested grouping survivability by day-1 (24 hrs) and day 2-4 (>24 hrs). The percent survivorship was tested between years (1984, 1985), sites (reference, Unit 2, Unit 3), and size-class (<100 mm, >100 mm), using a three-way analysis of variance weighted by fish number.

Impingement and return data for 1984 and 1985 were obtained from Southern California Edison's database. Statistical Analysis System (SAS) was used to perform all statistical analyses, particularly the procedures REG, GLM, SURVTEST, and LIFETEST (SAS 1983, 1985).

Table 1

Return efficiency of the fish return system at San Onofre Nuclear Generating Station, 1984. Fishes listed in order of abundance. Returns based on extrapolations from aliquot samplings.

Species	Total no. taken in	No. impinged	No. returned	% returned	Species	Total no. taken in	No. impinged	No. returned	% returned
Engraulis mordax	135,688	1,012	134,676	99.3	*Scomber japonicus*	16	4	12	75.0
Seriphus politus	50,566	6,197	44,369	87.7	*Girella nigricans*	13	0	13	100.0
Anchoa delicatissima	3,693	635	3,058	82.8	*Syngnathus* sp.	13	13	0	0
Atherinopsis californiensis	1,445	11	1,434	99.2	*Trachurus symmetricus*	13	2	11	84.6
Anchoa compressa	889	181	708	79.6	*Rhinobatos productus*	12	0	12	100.0
Xenistius californiensis	831	20	811	97.6	*Mustelus henlei*	10	0	10	100.0
Genyonemus lineatus	644	43	601	93.3	*Platyrhinoidis triseriata*	9	3	6	66.7
Atherinops affinis	601	11	590	98.2	*Torpedo californica*	9	2	7	77.8
Hermosilla azurea	436	0	436	100.0	*Cymatogaster aggregata*	8	3	5	62.5
Paralabrax clathratus	270	1	269	99.6	*Hypsoblennius* spp.	8	8	0	0
Peprilus simillimus	263	33	230	87.5	*Pleuronichthys ritteri*	8	4	4	50.0
Umbrina roncador	258	0	258	100.0	*Damalichthys vacca*	6	0	6	100.0
Hyperprosopon argenteum	227	6	221	97.4	*Hypsypops rubicundus*	6	0	6	100.0
Anisotremus davidsonii	211	1	210	99.5	*Citharichthys stigmaeus*	5	0	5	100.0
Porichthys notatus	111	108	3	2.7	*Scorpaena guttata*	5	3	2	40.0
Menticirrhus undulatus	108	1	107	99.1	*Sebastes paucispinis*	5	1	4	80.0
Paralabrax nebulifer	89	3	86	96.6	*Halichoeres semicinctus*	3	0	3	100.0
Phanerodon furcatus	80	21	59	73.7	*Heterodontus francisci*	3	0	3	100.0
Sardinops sagax	75	14	61	81.3	*Mustelus californicus*	3	1	2	66.7
Chromis punctipinnis	45	0	45	100.0	*Gymnura marmorata*	2	0	2	100.0
Paralichthys californicus	45	3	42	93.3	*Gibbonsia metzi*	1	1	0	0
Urolophus halleri	41	5	36	87.8	*Gibbonsia* sp.	1	1	0	0
Cheilotrema saturnum	38	0	38	100.0	*Hypsoblennius gilberti*	1	1	0	0
Roncador stearnsii	36	1	35	97.2	*Medialuna californiensis*	1	0	1	100.0
Leuresthes tenuis	35	10	25	71.4	*Otophidium scrippsi*	1	0	1	100.0
Atractoscion nobilis	25	0	25	100.0	*Pleuronichthys verticalis*	1	1	0	0
Embiotoca jacksoni	24	1	23	95.8	*Porichthys myriaster*	1	1	0	0
Heterostichus rostratus	20	18	2	10.0	*Squatina californica*	1	0	1	100.0
Sphyraena argentea	19	10	9	47.4	Total	196,978	8,395	188,583	95.7

Results

Diversion Efficiency

With some exceptions, fishes were diverted with high frequency into the FRS (Tables 1,2). This was particularly true in 1984, when 13 of the 15 most abundant species were diverted with 80% or better efficiency, 10 species exceeding 90%. The most abundant species, northern anchovy (*E. mordax*) and queenfish (*S. politus*), were diverted with 99.3% and 87.7% efficiency, respectively. The system worked particularly well for strong-swimming forms (i.e., jacksmelt, *Atherinopsis californiensis*; salema, *Xenistius californiensis*; topsmelt, *Atherinops affinis*; and kelp bass, *Paralabrax clathratus*). A high percentage of important sport and commercial fishes (*P. clathratus*; yellowfin croaker, *Umbrina roncador*; California corbina, *Menticirrhus undulatus*; California halibut, *Paralichthys californicus*) were also diverted with high frequency. The system did not appreciably divert weak-swimming species (such as plainfin midshipmen, *Porichthys notatus*; pipefish, *Syngnathus* spp.; giant kelpfish, *Heterostichus rostratus*).

Return rates of some species were not as high in 1985 as in the previous year (Tables 1,2). Nine of the 15 most abundant species were diverted 80% of the time or better. While *E. mordax* was diverted at about the same frequency between years (99.3 in 1984 vs. 94.3 in 1985), efficiencies for *S. politus* (87.7 vs. 73.7%), white croaker, *Genyonemus lineatus* (93.3 vs. 38.5%), deepbody anchovy, *Anchoa compressa* (79.6 vs. 49.4%), and slough anchovy, *Anchoa delicatissima* (82.8 vs. 4.5%) were considerably lower. With these exceptions, between-year return efficiency was similar.

No seasonal pattern was found in return efficiency based on an examination of the overall temporal pattern for seasonality by checking for significant autocorrelation. The three-way analysis of variance, with transformed return efficiency as the dependent variable and site, year, and size as the main effects, was then performed for the five species. We first analyzed samples collected on the same day between SONGS Unit 2 and 3. Using this matched data, it was found that *E. mordax* showed significant differences between year ($p < 0.05$), *S. politus* showed significant differences between size ($p < 0.001$) and year ($p < 0.001$), while the other three species analyzed showed no differences for any of the factors in the ANOVA.

Table 2

Return efficiency of the fish return system at San Onofre Nuclear Generating Station, 1985, based on operation criteria of 22-26 hours and four circulating pumps. Fishes listed in order of abundance. Returns based on extrapolations from aliquot sampling.

Species	Total no. taken in	No. impinged	No. returned	% returned
Engraulis mordax	210,108	11,951	198,157	94.3
Seriphus politus	104,394	27,431	76,963	73.7
Genyonemus lineatus	52,938	32,548	20,390	38.5
Anchoa delicatissima	27,514	26,284	1,230	4.5
Anchoa compressa	3,809	1,926	1,883	49.4
Umbrina roncador	2,026	5	2,021	99.8
Xenistius californiensis	1,885	61	1,824	96.8
Hyperprosopon argenteum	978	58	920	94.1
Phanerodon furcatus	819	280	539	65.8
Peprilus simillimus	488	88	400	82.0
Anisotremus davidsonii	284	2	282	99.3
Porichthys notatus	250	241	9	3.6
Menticirrhus undulatus	178	2	176	98.9
Atherinopsis californiensis	177	14	163	92.1
Paralabrax clathratus	165	4	161	97.6
Hermosilla azurea	158	0	158	100.0
Cymatogaster aggregata	145	90	55	37.9
Heterostichus rostratus	141	105	36	25.5
Atractoscion nobilis	138	18	120	87.0
Syngnathus species	116	116	0	0
Scomber japonicus	98	13	85	86.7
Paralichthys californicus	83	9	74	89.2
Urolophus halleri	74	4	70	94.6
Porichthys myriaster	69	61	8	11.6
Atherinops affinis	57	11	46	80.7
Sphyraena argentea	54	33	21	38.9
Paralabrax nebulifer	50	3	47	94.0
Trachurus symmetricus	46	3	43	93.5
Roncador stearnsii	42	2	40	95.2
Rhinobatos productus	39	1	38	97.4
Pleuronichthys ritteri	33	13	20	60.6
Embiotoca jacksoni	32	4	28	87.5
Xystreurys liolepis	32	18	14	43.8
Platyrhinoidis triseriata	30	20	10	33.3
Chromis punctipinnis	27	1	26	96.3
Leuresthes tenuis	27	19	8	29.6
Otophidium scrippsi	27	24	3	11.1
Torpedo californica	25	21	4	16.0
Myliobatis californica	21	1	20	95.2
Scorpaena guttata	19	10	9	47.4
Cheilotrema saturnum	17	0	17	100.0
Sebastes paucispinis	16	11	5	31.3
Mustelus californicus	13	4	9	69.2
Damalichthys vacca	12	1	11	91.7
Etrumeus teres	12	1	11	91.7
Medialuna californiensis	12	0	12	100.0
Citharichthys stigmaeus	7	7	0	0
Gymnura marmorata	7	3	4	57.1
Hypsoblennius spp.	7	7	0	0
Halichoeres semicinctus	5	1	4	80.0
Hypsopsetta guttulata	5	0	5	100.0
Mustelus henlei	5	3	2	40.0
Scorpaenichthys marmoratus	5	3	2	40.0
Triakis semifasciata	4	1	3	75.0
Gibbonsia sp.	3	3	0	0
Girella nigricans	3	0	3	100.0
Heterodontus franscisci	3	1	2	66.7
Leptocottus armatus	3	3	0	0
Rhacochilus toxotes	3	0	3	100.0
Oxyjulis californica	2	1	1	50.0
Sebastes serriceps	2	2	0	0
Synodus lucioceps	2	2	0	0
Amphistichus argenteus	1	1	0	0
Clupea harengus pallasii	1	0	1	100.0
Gibbonsia elegans	1	1	0	0
Hypsypops rubicundus	1	0	1	100.0
Microstomus pacificus	1	1	0	0
Paralabrax maculatofasciatus	1	0	1	100.0
Pleuronichthys verticalis	1	0	1	100.0
Sarda chiliensis	1	0	1	100.0
Sebastes auriculatus	1	1	0	0
Squalus acanthias	1	1	0	0
Symphus atricauda	1	1	0	0
Total	407,755	101,555	306,200	75.1

As no differences were found for any of the species in transformed return efficiency between the two units, analyses were performed using all sampling dates for the two units for the period 1984-85 in order to increase the sample size (detailed computations are presented in Appendix B). Analyses with all dates included showed significant differences for size and year for *E. mordax* (Fig. 6), where fish <100 mm were returned more efficiently than those greater ⩾100 mm, and the 1984 return efficiency was better than 1985 (Bonferroni test). Similarly, for *S. politus* the analyses revealed significant differences for size ($p < 0.001$). From the Bonferroni test and Figure 10, it is clear that the return efficiency in 1984 was consistently greater than 1985. In addition, return efficiency was better for fishes ⩾100 mm than for those <100 mm. For the other three species, *G. lineatus*, *H. argenteum*, and *P. furcatus* (Figs. 7-9), we again note that none of the factors showed significant differences. However, in examining their means it was seen that 1984 was consistently higher in return efficiency than 1985 and that lengths >100 mm was greater than <100 mm for all three species. It might be argued that the relatively few fish sampled over 100 mm would weaken the analysis of variance, but since no interactions were found in the ANOVA, this does not appear to be a problem.



Figure 6
Return efficiency (%) by length of northern anchovy, *Engraulis mordax*, at San Onofre Units 2 and 3.



Figure 8
Return efficiency (%) by length of walleye surfperch, *Hyperprosopon argenteum*, at San Onofre Units 2 and 3.



Figure 7
Return efficiency (%) by length of white croaker, *Genyonemus lineatus*, at San Onofre Units 2 and 3.



Figure 9
Return efficiency (%) by length of white surfperch, *Phanerodon furcatus*, at San Onofre Units 2 and 3.



Figure 10
Return efficiency (%) by length of queenfish, *Seriphus politus*, at San Onofre Units 2 and 3.

Survivorship

The results of the 96-hr survivorship studies are presented in Table 3. Individuals of most species survived both transit through the fish return system and 96 hrs in the holding net. There was little or no mortality for such species as *E. mordax*, *U. roncador*, and *X. californiensis*. No *Anchoa delicatissima* survived. With the exception of *E. mordax* and *S. politus*, all fishes survived in the reference net.

We examined most closely the survivorship of *E. mordax*, *S. politus*, and *G. lineatus*, three of the species given special attention in the return efficiency section. Survivorship of these three species, by length (10-mm increments) is shown in Figures 11-13. Figure 11 indicates that *E. mordax* small individuals (<60 mm) survived most frequently at both units. Mortality increased rapidly after 60 mm. Survivorship among *S. politus* exhibited a more complicated pattern (Fig. 12). Survivorship was poor among smaller fishes (<70 mm) traveling through Unit 2, and highest in 70-120 mm individuals. In contrast, smallest *S. politus* survived Unit 3 best and mortality increased with fish length. However, the ANOVA results indicate that this pattern probably only applies to day-1 survivorship. For *G. lineatus*, small individuals (≥70 mm) increased in survivorship with length at the two units (Fig. 13).

Table 3
Percent survivorship of species in 96-hour holding experiments, 1984-85. Numbers of individuals used in experiments are in parentheses.

	Experimental		
	Unit 2	Unit 3	Reference
Species	$n = 6$	$n = 8$	$n = 7$
Anchoa compressa	50.0 (2)		
Anchoa delicatissima	0.0 (95)		
Anisotremus davidsonii	100.0 (2)		100.0 (3)
Atherinopsis californiensis	100.0 (2)		
Atractoscion nobilis	100.0 (1)	100.0 (1)	
Chromis punctipinnis	100.0 (1)	100.0 (1)	
Cymatogaster aggregata	100.0 (1)		100.0 (4)
Damalichthys vacca		100.0 (4)	
Engraulis mordax	94.3(930)	97.9(4630)	14.8(108)
Genyonemus lineatus	49.5 (95)	25.0 (40)	100.0 (10)
Hermosilla azurea	100.0 (3)		
Heterostichus rostratus	100.0 (1)	100.0 (1)	
Hyperprosopon argenteum	100.0 (19)	100.0 (12)	100.0 (1)
Medialuna californiensis	100.0 (1)		
Menticirrhus undulatus		100.0 (1)	100.0 (2)
Paralabrax clathratus	100.0 (1)		100.0 (3)
Paralabrax nebulifer	100.0 (1)		100.0 (4)
Peprilus simillimus		100.0 (1)	
Phanerodon furcatus	100.0 (5)	94.7 (19)	100.0 (8)
Sebastes paucispinis		100.0 (1)	
Seriphus politus	31.6(753)	54.1 (846)	78.8(170)
Umbrina roncador	100.0 (58)	97.0 (133)	100.0 (15)
Xenistius californiensis	100.0 (21)	100.0 (38)	



Figure 11
Percent survivorship by length of northern anchovy, *Engraulis mordax*, in the reference net and San Onofre Units 2 and 3.



Figure 12
Percent survivorship by length of queenfish, *Seriphus politus*, in the reference net and San Onofre Units 2 and 3.



Figure 13
Percent survivorship by length of white croaker, *Genyonemus lineatus*, in the reference net and San Onofre Units 2 and 3.

Figure 14 presents the conditional probability of survival by reference experiments and Units 2 and 3 by size-class for the three species. Percent survivorship declined most precipitously by day-1, suggesting that survivorship through day-1 was most crucial. For this reason, analyses were conducted separating day-1 from days 2-4.

The three-way analysis of variance showed several statistically significant differences for *E. mordax* and *S. politus* and none for *G. lineatus* (Appendix C).

Site differences found in *E. mordax* for day-1 were due to the reference experiment being considerably lower in mean survivorship ($\bar{x}$ = 0.41) than the two units (unit 2, $\bar{x}$ = 0.97; unit 3, $\bar{x}$ = 0.94). No differences for *E. mordax* were found for days 2-4 for any effects.

Seriphus politus, however, exhibited interaction between year and site for both day-1 and day 2-4 analyses when averaged over size. Examination of the day-1 mean percent survivorship showed that fishes in Unit 3 had better survivorship in 1984 than 1985. This was not the case at Unit 2. Here, in contrast, *S. politus* survivorship increased in 1985 (Appendix D). Thus, though there appears to be a between-site difference in *S. politus* survivorship, this Anova interaction makes it impossible to delineate whether the site differences were truly significant. Examination of days 2-4 mean percent survivorship showed that Unit 3 survivorship decreased relatively more than both reference and Unit 2 in 1985, causing significant interactions. A main-effect difference also existed for year. Main-effect differences for the year are due to 1984 survivorship percentage exceeding that of 1985.

Potentially, there may be several other variables that might explain differences seen in *S. politus* survivorship. The total biomass of fishes in the return sample was thus used as a covariate of percent survivorship that may eliminate the interaction seen. The three-way analysis of variance was therefore repeated, using biomass as a covariate for day-1 and days 2-4 survivorship for *S. politus* only. However, the results suggested that total biomass of fishes was an insignificant covariate for percent survivorship of *S. politus*.

Discussion

Most species of fish entrapped by the Units 2 and 3 intakes were diverted efficiently by the SONGS fish return system. This was particularly true of such species as *Paralabrax clathratus*, *Xenistius californiensis*, *Umbrina roncador*, *Engraulis mordax*, and *Seriphus politus*. Very poor swimmers (i.e., *Syngnathus* spp. and *Heterostichus rostratus*) were not diverted well.

To a certain extent, these results were influenced by fish size. For instance, diversion, highest among small *E. mordax*, gradually declined with fish length. However, this pattern was the exception. In most species, larger individuals were diverted with highest frequency. In a similar study from a freshwater system, Lawler et al. (1982b) found that larger species and larger individuals within each species displayed the highest diversion efficiency. It is noteworthy that *E. mordax* did not follow this pattern, as the design of this system



Figure 14
Conditional probability of survival of *Engraulis mordax*, *Seriphus politus*, and *Genyonemus lineatus* in the reference net and San Onofre Units 2 and 3 experiments.

(Schuler and Larson 1975) was derived from modeling studies using adult *E. mordax* as the principal species for fine-tuning the design of various system components.

The potential ability of fish in the smallest length-classes (<50 mm) to pass through the 9.5-mm mesh of the traveling screens and impingement baskets may confound our evaluation of diversion efficiency of larval and juvenile fishes. Frequently, large schools of larval and juvenile *E. mordax* appeared in fish return samples while being totally absent in the paired impingement samples. This absence may be attributed to the inability of the traveling screens to adequately filter the entrained water of fish of this size. Any fish of this size that was impinged would most likely be washed through the mesh of the impingement basket by the vigorous action of the sluicing water. We have observed this to occur at Unit 2, where larval anchovy are occasionally present in the water surrounding the impingement basket. Very small *E. mordax* appear especially susceptible because their small lateral surface area offers little "impingible" surface when compared with deeper-bodied fish of the same length, such as *S. politus*. Thus, the high diversion efficiency of small *E. mordax*, compared with *S. politus*, may have been due to the ability of small *E. mordax* to pass through the traveling screens and impingement baskets. This would bias our samples toward those small anchovies which were diverted and hence counted.

There was a lower rate of return for some species in 1985 compared with 1984. The phenomenon does not appear to be related to differences in fish sizes between years, as return rates for all size-classes of *S. politus* and nearly all of *E. mordax* were lower in 1985. As one examines the information for the two years, we see that the number of fish were higher in 1985, yet overall return efficiency was lower for 1985. There is a possibility that return efficiency may be negatively related to fish density. Examination of fish density patterns are beyond the scope of this paper although it is of interest biologically. Future studies might consider fish density when examining return efficiency.

Generally, fishes were not only diverted through the fish return system, but also survived the experience (based on our holding-net experiments). However, some species fared better than others. *Engraulis mordax*, *U. roncador*, and *X. californiensis* rarely died, while *Anchoa delicatissima* always did.

Most fishes survived in the reference nets. However, high mortality occurred in *E. mordax* and *S. politus*. We believe, at least in the case of *E. mordax*, this was due to the species' apparent inability to school well in numbers less than about 40, a number we were unable to capture at any one time.

Our diver observations of captured fish in the reference nets indicate that small numbers of these two species frequently display difficulty in forming schools, show erratic swimming behavior, and appear to contact the net more often than individuals within large schools. Obligate schooling fishes, such as queenfish and anchovy, rely on visual cues to provide the attractive force between members of the school (Hemmings 1966, Partridge 1982), and within the school each member expends less energy than if swimming alone (Breder 1976). The small numbers of individuals in the reference nets may deter captured fish from forming cohesive schools. Any resultant increase in energy expenditure or collisions with the net would certainly have a deleterious influence on the survivorship of these relatively fragile species.

We noted some size-specific mortality, in *E. mordax*. Within this species mortality increased with size. Based on Figure 13, it would appear that there is a between-unit difference in size-specific *S. politus* mortality. However, based on our analysis of variance, this pattern was seen for day-1 but not for days 2-4.

We also noted differences in survivorship of *S. politus* between the two units in 1984 and 1985. For day-1 in 1984, survivorship was lower at Unit 2, but was lower in 1985 at Unit 3. This occurrence is not readily explainable, even when attempts were made to use total fish biomass as a covariate.

In some cases, variable survivorship of juveniles in the holding experiments may reflect the inability of some small individuals to orient to the discharge flow within the return conduit. Diver observations indicate that returned queenfish less than approximately 60 mm in length show a large variation in their ability to orient to the conduit flow. As they exit the discharge, most appear to "tumble" out of the conduit instead of slowly backing down with the flow, as larger individuals and members of other species have been observed to do. Tumbling within the return conduit would certainly increase the susceptibility of fish to abrasion and subsequent descaling and may subject them to fast, potentially damaging changes in hydrostatic pressure. Both of these effects have been demonstrated to cause mortalities in *S. politus* (Johnson 1981). In addition, we have observed that small queenfish often appear initially stunned or disoriented upon discharge from the conduit and are less able to school effectively within the holding net. These individuals frequently collide with the net walls and thereby may further increase their trauma.

Factors we cannot control or test for may also be important in determining the survival of fish discharged from the return system. Some of the fishes which survive passage through the plant are eaten by predators upon discharge from the return system. Small groups of barred sand bass (*Paralabrax nebulifer*) and kelp bass (*Paralabrax clathratus*) and solitary California halibut (*Paralichthys californicus*) congregate near the discharge, having apparently associated the conduit opening with food. However, it is the infrequent visits of schooling predators such as jack mackerel (*Trachurus symmetricus*), Pacific mackerel (*Scomber japonicus*), and large *S. politus* that appear to result in the highest predation pressure. We observed schools of these predators (as well as those of California barracuda, *Sphyraena argentea*) on 13 of 80 days of observations on the return system's discharge. Although we have observed this behavior only occasionally during the day, we do not know how often nocturnal or crepuscular predators frequent the fish discharge. Information such as this may be important in determining the time and frequency of running the fish return system to insure maximum survival of return fish.

We should also mention that survivorship in the net is "relative" survivorship. It is likely lower than true survivorship (excluding predation effects), due to such factors as the trauma to fish thrust into the holding net and restricted foraging of the individuals.

Although unexplained variations in efficiency of survivorship made the analysis of the effectiveness of the SONGS fish return system complex, we feel that these results strongly support the success of the SONGS fish return system. There is no doubt that this system is a considerable improvement in fish conservation over previously applied technologies.

Acknowledgments

This work was funded by Southern California Edison Contract C0206906. We wish to thank John Palmer and Kevin Herbinson, SCE, for their support. Mike Curtis, Craig Heberer, and Walter Marsh also provided considerable assistance. Edward DeMartini, Keith Parker, and James Bence gave us extensive critical analyses of this manuscript. Sara Warschaw and Susan Muth typed this paper.

Citations

Bailey, J.R.

1977 Tables of the Bonferroni T Statistic. J. Am. Stat. Assoc. 72:469-478.

Breder, C.M.

1976 Fish schools as operational structures. Fish. Bull., U.S. 74:471-502.

DeMartini, E.E., L. Bost, R.K. Fountain, E. Koehrn, and E. Roberts

1984 Preliminary evaluation of impingement and diversion of fishes at SONGS Unit 2; update of lampara seine monitoring data for coastal pelagic fishes; and update of otter trawl data for benthic soft-bottom fishes, p. 4-5. Trimestral Rep. UCSB Fish Proj., Mar. Sci. Inst., Univ. Calif., Santa Barbara, CA 93106.

Downs, D.I., and K.R. Meddock

1974 Design of fish conserving intake system. Proc. Am. Soc. Civil Eng. 100:191-205.

EPRI (Electric Power Research Institute)

1981 Impingement and entrainment: An updated annotated bibliography. EPRI Tech. Rep. EA-1855, prepared by Atomic Industrial Forum, Inc., and Oak Ridge Natl. Lab. (F.E. Yost and M.S. Uziel, principal investigators) for Electric Power Res. Inst., 3412 Hillview Ave., Palo Alto, CA 94304, 317 p.

1984 Advanced intake technologies study. EPRI Tech. Rep. CS-3644, prepared by Stone and Webster Eng. Corp. (Y.G. Mussalli, proj. eng.) for Electric Power Res. Inst., 3412 Hillview Ave., Palo Alto, CA 94304, 276 p.

Helvey, M.

1985 Behavioral factors influencing fish entrapment at offshore cooling-water intake structures in Southern California. Mar. Fish. Rev. 47(1):18-26.

1987 Selective removal of reef fishes associated with offshore cooling-intake structures. J. Appl. Ecol. 24:1-12.

Hemmings, C.C.

1966 Olfaction and vision in fish schooling. J. Exp. Biol. 45:449-464.

Johnson, L.

1981 Report to Southern California Edison, Research and Development, May 1981, Rosemead, CA 91770.

Lawler, Matusky and Skelly Engineers

1982a Intake technology review. Prepared for Southern California Edison, Research and Development Ser. 82-RD-164. Lawler, Matusky and Skelly, 1 Blue Hill Plaza, Pearl River, NY 10965.

1982b Evaluation of the angled screen fish diversion system at Oswego Steam Station Unit 6. Interim Rep. to Niagara Mohawk Power Corp., Syracuse, NY, 68 p. Lawler, Matusky and Skelly, 1 Blue Hill Plaza, Pearl River, NY 10965.

Lorda, E., and S.B. Saila

1986 A statistical technique for analysis of environmental data containing periodic variance components. Ecol. Model. 32:59-69.

Partridge, B.L.

1982 The structure and function of fish schools. Sci. Am. 246:114-123.

SAS

1983 Supplemental Library Users Guide (SUGI). SAS Inst., Cary, NC 27511.

1985 User's Guide Statistics, Version 5 Edition. SAS Inst., Cary, NC 27511, 956 p.

Schuler, V.J., and L.E. Larson

1975 Improved fish protection at intake systems. Proc. Am. Soc. Civil Eng. 101:897-910.

Yuge, J.E., and K.T. Herbinson

1985 Study introduction and generating station description. *In* Report on 1984 data. Marine Environmental Analysis and Interpretation, San Onofre Nuclear Generating Station, p. 1-10. Rep. 85-RD-37, Southern California Edison Co., Rosemead, CA 91770.

Appendix A

Actual size-measurement data for five species impinged and entrained at Units 2 and 3, San Onofre Nuclear Generating Station.

Size (in 10-mm increments)	*Engraulis mordax*	*Genyonemus lineatus*	*Hyperprosopon argenteum*	*Phanerodon furcatus*	*Seriphus politus*
1984					
10	1	·	·	·	·
20	30	·	·	·	42
30	1,053	1	·	·	526
40	1,511	6	3	9	1,921
50	1,211	15	3	15	2,204
60	582	18	3	4	1,719
70	223	13	15	1	1,657
80	94	41	8	1	882
90	35	14	3	·	305
100	7	3	6	1	191
110	3	1	5	·	329
120	·	·	2	1	248
130	·	·	·	·	127
140	·	·	1	·	56
150	·	1	·	1	16
160	·	2	·	2	8
170	·	1	·	·	6
180	·	·	·	·	2
1985					
10	·	·	·	·	2
20	161	1	·	·	74
30	1,630	36	·	·	837
40	2,828	507	1	8	2,954
50	2,037	2,590	27	247	2,875
60	1,130	2,187	96	79	1,897
70	667	482	80	35	1,395
80	251	89	53	16	1,777
90	94	20	18	3	2,198
100	151	7	7	·	2,332
110	79	·	·	·	2,016
120	33	1	·	·	1,458
130	9	·	·	·	959
140	1	·	·	·	486
150	·	1	·	·	246
160	·	·	·	2	89
170	·	3	·	3	42
180	·	1	·	1	6
190	·	2	·	2	9
200	·	·	·	·	2
210	·	1	·	·	·
230	·	1	·	·	·

Appendix B

Three-way ANOVA of *Engraulis mordax*, *Genyonemus lineatus*, *Hyperprosopon argenteum*, *Phanerodon furcatus*, and *Seriphus politus* for transformed return efficiency using matched dates (n = 37) versus all criteria sampling dates for 1984-85. Variable: Ln(1% return).

Species:	*Engraulis mordax*			*Genyonemus lineatus*			*Hyperprosopon argenteum*			*Phanerodon furcatus*			*Seriphus politus*		
	Type II			Type II			Type II			Type II			Type II		
Factor	*DF*	*SS*	*P*-value	*DF*	*SS*	*P*-value	*DF*	*SS*	*P*-value	*DF*	*SS*	*P*-value	*DF*	*SS*	*P*-value
Matched dates															
Model	7	79.76	0.13	6	36.62	0.50	5	49.86	0.25	5	48.30	0.31	7	153.21	0.001
Error	81	496.45	—	51	346.47	—	33	236.47	—	25	190.95	—	123	344.40	—
Year	1	39.35	0.02	1	0.49	0.79	1	19.82	0.11	1	0.01	0.98	1	52.57	0.0001
Size	1	18.18	0.11	1	24.22	0.06	1	3.11	0.51	1	18.85	0.13	1	98.79	0.0001
Site	1	8.03	0.28	1	7.20	0.31	1	0.15	0.89	1	0.11	0.90	1	3.18	0.29
Year*Size	1	0.01	0.98	1	1.30	0.66	*			1	8.67	0.30	1	5.68	0.16
Year*Site	1	9.53	0.24	1	3.51	0.47	1	0.42	0.81	1	4.15	0.47	1	0.11	0.84
Size*Site	1	0.00	0.98	1	0.79	0.73	1	0.16	0.80	*			1	0.32	0.74
Year*Size*Size	1	0.26	0.84	*			*			*			1	0.02	0.93
All dates															
Model	7	113.68	0.03	7	100.20	0.08	7	45.97	0.36	6	35.47	0.66	7	136.58	0.0001
Error	130	927.92	—	90	672.57	—	58	339.97	—	40	343.92	—	217	716.54	—
Year	1	36.34	0.03	1	13.73	0.18	1	8.46	0.23	1	0.88	0.75	1	40.22	0.0006
Size	1	27.62	0.05	1	26.94	0.06	1	13.69	0.13	1	25.25	0.09	1	92.66	0.0001
Site	1	2.57	0.55	1	13.10	0.19	1	3.03	0.47	1	1.21	0.71	1	1.91	0.45
Year*Size	1	15.62	0.14	1	0.03	0.95	1	1.26	0.64	1	2.13	0.62	1	1.27	0.53
Year*Site	1	10.90	0.22	1	0.39	0.82	1	4.51	0.38	1	0.41	0.83	1	0.70	0.64
Size*Site	1	4.35	0.44	1	28.01	0.06	1	0.81	0.71	1	7.24	0.36	1	0.25	0.78
Year*Size*Size	1	0.08	0.92	1	4.42	0.44	1	0.62	0.74	*			1	3.47	0.31

*Due to zero counts of total abundances at least once during the study, the interaction could not be considered in the model.

Appendix C

Three-way ANOVA of *Engraulis mordax*, *Genyonemus lineatus*, and *Seriphus politus* for percent survivorship in the 96-hour study.

Species:	*Engraulis mordax*			*Genyonemus lineatus*			*Seriphus politus*		
	Type II			Type II			Type II		
	DF	*SS*	*P*-value	*DF*	*SS*	*P*-value	*DF*	*SS*	*P*-value
Percent survivorship at end of day-1									
Model	5	23.90	0.16	6	1.83	0.57	9	104.60	0.01
Error	4	6.55	—	2	0.60	—	29	52.96	—
Year	1	0.28	0.70	1	0.58	0.30	1	0.62	0.56
Size	1	0.07	0.84	1	0.00	1.00	1	0.13	0.79
Site	2	23.56	0.05	2	0.15	0.80	2	33.71	0.01
Year*Size	1	0.11	0.81	*			1	4.44	0.13
Year*Site	*			2	0.04	0.93	2	41.14	0.01
Size*Site	*			*			2	9.19	0.10
Percent survivorship from day-2 to end of study									
Model	5	13.11	0.36	6	10.14	0.50	9	43.83	0.01
Error	2	2.52	—	1	0.87	—	23	18.18	—
Year	1	0.03	0.90	1	4.33	0.27	1	22.07	0.01
Size	1	1.65	0.37	1	0.00	1.00	1	0.05	0.81
Site	2	11.25	0.18	2	6.71	0.34	2	7.99	0.02
Year*Size	1	0.06	0.85	*			1	0.59	0.39
Year*Site	*			2	0.68	0.75	2	13.93	0.01
Size*Site	*			*			2	0.43	0.76

*Due to zero counts of fish at least once during the study, the interaction could not be considered in the model.



Appendix D

Significant interaction plots of site*year for *Seriphus politus* in the 96-hour study.

NOAA TECHNICAL REPORT NMFS

Guidelines for Contributors

FORM OF MANUSCRIPT

Authors are encouraged to retain manuscripts on word processing storage media, such as diskettes, floppy disks, cassette or magnetic tapes, and submit a hard copy run from the storage media. Hardcopy should be typed double-spaced on white bond paper in the following sequence:

Title page
Contents
Abstract
Text
Acknowledgments
Citations
Text footnotes
Appendix
Tables
Figure Legends
Figures

CONTENTS OF MANUSCRIPT

Contents. Contains all text headings with page numbers omitted.

Abstract. Not to exceed one double-spaced typed page. Should include a sentence or two explaining to the general reader why the research was undertaken and why the results should be viewed as important. Abstract should convey the main point of the paper and outline the results or conclusions. Footnotes and references do not belong in the abstract.

Text. A brief introduction should portray the broad significance of the paper. The entire text should be intelligible to readers from different disciplines. All technical terms should be defined. Follow the *U.S. Government Printing Office Style Manual*, 1984 edition.

Fishery nomenclature follows the American Fisheries Society Special Publication No. 12, *A List of Common and Scientific Names of Fishes from the United States and Canada*, 4th edition, 1980.

Abbreviations and symbols. Define all symbols, abbreviations, and acronyms. Define all symbols in equations and formulas. Abbreviate units of measure only when used with numerals or in tables and figures where there is lack of space. Periods are rarely used except for et al., vs., e.g., i.e., etc.

Measurements should be expressed in metric units. Other equivalent units may be given in parentheses.

Text footnotes. Type on a separate sheet from the text. Footnotes are not used for reference material or personal communications, but rather to explain or define terms in the text and for contribution numbers.

Personal communications are noted in parentheses in the text (name, affiliation, brief address including zip code, month and year).

Citations. All written sources should be listed in the Citations section, including unpublished and processed material. In text, cite as Smith and Jones (1977) or (Smith and Jones 1977); if more than one citation, list chronologically (Smith 1936; Jones 1975; Doe 1977). All sources cited in the text should be listed alphabetically by the senior authors' surname under the heading CITATIONS. Abbreviations of periodicals and serials should conform to the Biosis Data Base™. The author is responsible for the accuracy of all citations.

Acknowledgments. Gather all acknowledgments into a brief statement at the end of the text. Give credit only for exceptional contributions and not to those whose contributions are part of their normal duties.

Tables. Tables should supplement, not duplicate, the text. Each table should appear on a separate page, and be numbered consecutively. Headings should be short but amply descriptive so that the reader need not refer to the text. For values less than 1, zeros should precede all decimal points. Give each column a heading with units of measure indicated in parentheses. Do not change the unit of measure within a column. Table footnotes should be numbered consecutively in Arabic numerals across the page from left to right and down; to avoid confusion with exponents, place them to the left of numerals, e.g., $^{3}10$. All tables should be cited consecutively in the text and their placement, where first mentioned, indicated in the left-hand margin of the manuscript page. Lengthy tables that are typed in the proper format and are clean and legible can be used camera-ready.

Figures. Photographs and line drawings should be of professional quality—clear and concise—and reducible to 42 picas for full-page width or to 20 picas for a single-column width, and a maximum of 57 picas high. All illustrations must be proportioned so that they are legible when reduced to fit the page format. Line weight and lettering should be sharp and even. Lettering on graphs should be upper and lower case, and vertical lettering should be avoided as much as possible (except for vertical, y, axis). For values less than 1, zeros should precede all decimal points. Reproductions of line art work are accepted in the form of photographic prints from negatives or photomechanical transfer. Halftones should be sharply focused with good contrast. Micron rules should be inserted on electron micrographs, even when magnification is included in the figure legend. There should be good distinction between identifying letters (preferably press-on) and background of photograph. All figures should be cited consecutively in the text and their placement, where first mentioned, indicated in the left-hand margin of the manuscript page. Each figure should be labeled in pencil on the back. DO NOT SEND original or photographic prints of figures to the Scientific Editor; they will be requested when the manuscript is accepted for publication.

Copyright. Government publications are in the public domain, i.e., they are not protected by copyright.

SUBMISSION OF MANUSCRIPTS

Send original hardcopy and two duplicated copies of the manuscript to:

Dr. Andrew E. Dizon, Scientific Editor
NOAA Technical Reports NMFS
Southwest Fisheries Center
National Marine Fisheries Service, NOAA
P.O. Box 271
La Jolla, CA 92038-0271

Indicate whether word processing storage media are available, but do not forward disks until requested by Scientific Editor or Scientific Publications office.

Copies of published reports. For single monographs, the senior author will be provided with 50 copies free of charge and 50 to his or her organization. For collections of papers, each senior author will receive 50 separates and one cover, and his or her organization will receive the same. Additional copies may be purchased in lots of 100.

ADDITIONAL INFORMATION

Questions regarding editorial policy and procedures may be forwarded to:

Nancy Peacock, Managing Editor
NOAA Technical Reports NMFS
National Marine Fisheries Service, NOAA
Scientific Publications Office
Bin C15700
Seattle, WA 98115

UNITED STATES
DEPARTMENT OF COMMERCE
NATIONAL OCEANIC AND ATMOSPHERIC
ADMINISTRATION
NATIONAL MARINE FISHERIES SERVICE
SCIENTIFIC PUBLICATIONS OFFICE
BIN C15700
SEATTLE, WA 98115
OFFICIAL BUSINESS
Penalty for Private Use, $300

BULK RATE
POSTAGE & FEES PAID
U.S. Department of Commerce
Permit No. G-19

NOAA SCIENTIFIC AND TECHNICAL PUBLICATIONS

The National Oceanic and Atmospheric Administration was established as part of the Department of Commerce on October 13, 1970. The mission responsibilities of NOAA are to assess the socioeconomic impact of natural and technological changes in the environment and to monitor and predict the state of the solid Earth, the oceans and their living resources, the atmosphere, and the space environment of the Earth.

The major components of NOAA regularly produce various types of scientific and technical information in the following kinds of publications:

PROFESSIONAL PAPERS—Important definitive research results, major techniques, and special investigations.

CONTRACT AND GRANT REPORTS—Reports prepared by contractors or grantees under NOAA sponsorship.

ATLAS—Presentation of analyzed data generally in the form of maps showing distribution of rainfall, chemical and physical conditions of oceans and atmosphere, distribution of fishes and marine mammals, ionospheric conditions, etc.

TECHNICAL SERVICE PUBLICATIONS—Reports containing data, observations, instructions, etc. A partial listing includes data serials; predictions and outlook periodicals; technical manuals, training papers, planning reports, and information serials; and miscellaneous technical publications.

TECHNICAL REPORTS—Journal quality with extensive details, mathematical developments, or data listings.

TECHNICAL MEMORANDUMS—Reports of preliminary, partial, or negative research or technology results, interim instructions, and the like.

Information on availability of NOAA publications can be obtained from:

U.S. Department of Commerce
National Technical Information Service
5285 Port Royal Road
Springfield, VA 22161